

Preferences of People with Hereditary Angioedema for On-Demand Treatment: A US-Based Qualitative Study

Don Bukstein,¹ Vibha Desai,² Ledia Goga,^{2a} Shawn Czado,² Michelle Brown,³ Kelley Myers,³ Paul Audhya,² Russell A. Settignano,⁴ Paula Busse⁵

¹The PBL Institute, Madison, Wisconsin, USA; ²KalVista Pharmaceuticals, Inc., Cambridge, Massachusetts, USA; ³RTI International, Inc., North Carolina, USA; ⁴Brown University, Providence, Rhode Island, USA; ⁵Department of Medicine, Division of Clinical Immunology, Mount Sinai, New York, USA.

^aEmployee of KalVista Pharmaceuticals at the time the study was conducted

Background

- Hereditary angioedema (HAE), a rare genetic disorder, is characterized by recurrent and unpredictable episodes of subcutaneous or submucosal swelling which can affect the abdomen, extremities, genitals, face, and larynx¹
- All currently approved HAE on-demand treatments must be administered parenterally, which results in significant treatment burden

Objective

- The objectives of this qualitative study were to understand patients' likes and dislikes related to their current on-demand treatment, their attack experiences and route of administration (ROA) preferences for on-demand treatment

Methods

- The US Hereditary Angioedema Association (HAEA) recruited 20 people living with type 1 or type 2 HAE to be interviewed
- Participants were not informed of the identity of the study sponsor
- Study population included both adults (18 to 69yrs) and adolescents (12 to 17yrs); has had at least one HAE attack within the past six months; currently taking on-demand treatment (C1-INH replacement or bradykinin receptor B2 antagonist or kallikrein inhibitor)
- The sampling plan aimed to obtain half of each age group currently taking both on-demand treatment and long-term prophylaxis (LTP) and half taking only on-demand treatment
- Open-ended questions were asked to participants to understand their likes and dislikes associated with their current on-demand treatment
- Open-ended questions were then asked to understand the trade-offs patients are willing to make when choosing a preferred ROA. Hypothetical self-administered injection and oral on-demand treatments were initially presented with similar efficacy and tolerability/mild side-effect risk profiles (Figure 1), which were then made better/worse depending upon participants' initial treatment choice
- Profiles were based on US package inserts of on-demand injection treatments and clinical trial data for oral on-demand treatment in development

Figure 1. Hypothetical Trade-off Scenario

Features	Treatment A	Treatment B
How you take the treatment	Self-injection administered at home	Pill
How long it takes to feel a little improvement in your symptoms?	Same	Same
Risk of mild to moderate pain after the injection	Yes, more than 9 out of 10 people	No
Increased risk of additional mild side effects (e.g., nausea, dizziness and headache)	Yes, fewer than 1 out of 10 people	Yes, fewer than 1 out of 10 people
Which option would you choose?	Treatment A <input type="checkbox"/>	Treatment B <input type="checkbox"/>

Table 1. Respondent Characteristics

Characteristic	Adolescents (n = 10)	Adults (n = 10)	Total (N = 20)
Age, mean years (SD) [min-max]	15.5 (1.5) [12-17]	36.7 (16) [18-60]	26.1 (8.9) [12-60]
Gender, n (%)			
Female	5 (50)	6 (60)	11 (55)
Male	5 (50)	4 (40)	9 (45)
Race/ethnicity, n (%)			
African American or Black	1 (10)	1 (10)	2 (10)
Hispanic, Latin American, or Latinx	4 (30)	—	4 (20)
Middle Eastern or North African	—	1 (10)	1 (5)
White	5 (50)	8 (80)	13 (65)
Age at HAE diagnosis, mean years (SD)	6.7 (4.3)	17.2 (12)	11.9 (8.3)
HAE type 1, n (%)	10 (100)	7 (70)	17 (85)
Number of attacks, last 6 months, n (SD)	4.4 (5.1)	4.5 (5.0)	4.5 (5.0)
Most recent attack location, n (%)			
Face	1 (10)	—	1 (5.0)
Extremities	3 (30)	4 (40)	7 (35)
Abdomen	5 (50)	4 (40)	9 (45)
Throat	1 (10)	2 (20)	3 (15)
Lifetime specific attack location, n (%)			
Abdominal	8 (80)	9 (90)	17 (85)
Throat	4 (40)	6 (60)	10 (50)
Current type of HAE treatment, n (%)			
On-demand treatment and LTPT	8 (80) ^c	5 (50)	13 (65)
On-demand treatment only	2 (20)	5 (50)	7 (35)
On-demand treatment used for most recent attack, n (%)			
Firazyr, icatibant ^a	1 (10)	8 (80)	9 (45)
Beriner ^b	3 (30)	1 (10)	4 (20)
Ruconest, conestat alfa	3 (30)	—	3 (15)
Used LTPT ^b	3 (30)	1 (10)	4 (20)

LTPT = long-term prophylactic treatment; SD = standard deviation.

^a While indicated for individuals aged 18 years and older, 1 adolescent (aged 15 years) reported recent first-time use of Firazyr for their on-demand treatment.

^b One adult (Haegarda) and 3 adolescents (Haegarda, Orladeyo, Takhzyro) reported using their LTPT as an on-demand treatment for their most recent attack. At screening, these participants reported use of Firazyr (n = 1) and Beriner (n = 3) as their current on-demand treatment.

^c Although the sampling plan aimed to obtain half of each age group currently taking both on-demand treatment and LTPT and half taking only on-demand treatment, this was not able to be achieved in the adolescent cohort.

Table 3. Patient Preferences

	Adolescents (n = 10)	Adults (n = 10)	Total (N = 20)
Treatment choice, n (%)			
Treatment A (self-administered injection)	—	—	—
Treatment B (oral)	10 (100)	10 (100)	20 (100)
Reasons for choosing treatment B, n (%)^a			
Less pain/burning	5 (30)	5 (50)	10 (50)
Convenient to take/carry	3 (30)	4 (40)	7 (35)
Only 4 hours required before second dose	4 (40)	2 (20)	6 (30)
Safer (due to no injection/infusion)	1 (10)	—	1 (5)
No needles	1 (10)	1 (10)	2 (10)
Less time to take	2 (20)	1 (10)	3 (15)
Responses to trade-off scenarios			
Treatment A offered "substantial improvement" (vs. Treatment B, "little improvement") within the same timeframe, n (%)			
Treatment A choice	9 (90)	8 (80)	17 (85)
Treatment B choice	1 (10)	2 (20)	3 (15)
The risk for mild side effects was higher for Treatment B?			
Odds before switching to Treatment A, n (%)^b			
< 5 in 10	2 (20)	3 (30)	5 (25)
≥ 5 in 10	8 (80)	7 (70)	15 (75)

SD = standard deviation.

^a Multiple response question; percentages sum to greater than 100% per column.

^b When 14 participants were asked about the specific side effects of headache, nausea, and dizziness, 9 participants reported that they were more likely to tolerate a headache; 5 participants reported that they were less likely to tolerate a headache.

Disclosures

This study was sponsored by KalVista Pharmaceuticals, Inc. All authors met the ICMJE authorship criteria and had full access to relevant data. The authors had full editorial control of the data presented and provided final approval of all content. Neither honoraria nor payments were made for authorship.

Don Bukstein – Reports being a speaker for Regeneron.

Vibha Desai, Ledia Goga, Shawn Czado and Paul Audhya – Employees of KalVista Pharmaceuticals, Inc. (*at time of the publication development)

Michelle Brown and Kelley Myers – Employees of RTI-Health Solutions.

Russell A. Settignano – Advisor for Biocryst, CSL Behring, and KalVista Pharmaceuticals, Inc., speaker for Biocryst, Pharming and Takeda, and an independent contractor for Biocryst and KalVista Pharmaceuticals, Inc.

Paula Busse – Advisor/Consultant and/or Grant/Research Support: CSL Behring, KalVista Pharmaceuticals, Inc., Biocryst Pharmaceuticals, Takeda, Regeneron, and Novartis.

Results

Table 2. Reported "Likes" and "Dislikes" of Most Recent Acute Attack Treatment by Mode of Administration^a

Characteristic	Adolescents ^b , n (%)				Adults ^c , n (%)			Total, n (%)			
	SCI (n = 3)	IVI (n = 6)	Pill (n = 1)	Total (n = 10)	SCI (n = 9)	IVI (n = 1)	Total (n = 10)	SCI (n = 12)	IVI (n = 7)	Pill (n = 1)	Total (N = 20)
Likes^d, n (%)											
Effective/reliable; "it works"	—	4 (67)	1 (100)	5 (50)	8 (89)	—	8 (80)	8 (80)	4 (40)	1 (100)	13 (65)
Feeling of the medicine going in (emotional relief)	—	3 (50)	—	3 (30)	—	—	—	—	3 (30)	—	3 (15)
Easy to inject (intravenous and subcutaneous)	—	1 (17)	—	1 (10)	1 (11)	—	1 (10)	1 (10)	1 (10)	—	2 (10)
Familiar/comfortable	—	1 (17)	—	1 (10)	—	—	—	—	1 (10)	—	1 (5)
Easy to constitute (e.g., referenced previous more cumbersome process)	—	1 (17)	—	1 (10)	—	—	—	—	1 (10)	—	1 (5)
SCI (vs. IVI)	1 (10)	—	—	1 (10)	1 (11)	—	1 (10)	1 (10)	1 (10)	—	2 (10)
Cost (e.g., "affordable")	—	—	—	—	1 (11)	—	1 (10)	1 (10)	—	—	1 (5)
Portable	—	—	1 (100)	1 (10)	2 (22)	—	2 (20)	2 (20)	—	1 (100)	3 (15)
Dislikes^d, n (%)											
Painful/burning injection	1 (33)	3 (50)	—	4 (40)	3 (33)	—	3 (30)	4 (40)	3 (30)	—	7 (35)
Takes too long to work (efficacy)	1 (33)	1 (17)	—	2 (20)	3 (33)	—	3 (30)	4 (40)	1 (10)	—	5 (25)
Cannot easily take with you (refrigeration and travel)	1 (33)	—	—	1 (10)	3 (33)	—	3 (30)	4 (40)	—	—	4 (20)
Needles/injections (fear/avoidance)	—	—	—	—	3 (33)	—	3 (30)	3 (30)	—	—	3 (15)
Burden/hassle of administration (i.e., time commitment; refrigeration; preparation)	—	2 (33)	—	2 (20)	2 (22)	1 (100)	3 (30)	2 (20)	3 (30)	—	5 (25)
Dependent on others for administration	—	2 (33)	—	2 (20)	—	—	—	—	2 (20)	—	2 (10)
Body-weight sensitive (i.e., 1 participant was administered too low of a dose due to a recent weight gain)	—	1 (17)	—	1 (10)	1 (11)	—	1 (10)	1 (10)	1 (10)	—	2 (10)
Same infusion site (e.g., they would like to be able to administer in other places)	1 (33)	1 (17)	—	2 (20)	—	—	—	1 (10)	1 (10)	—	2 (10)
High cost	—	—	—	—	1 (11)	—	1 (10)	1 (11)	—	—	1 (5)

IVI = intravenous; IVI = intravenous infusion; SCI = subcutaneous injection.

^a Reported likes and dislikes were based on the treatment used for their most recent attack, including 4 participants who used a long-term prophylactic treatment.

^b Ten adolescents took Firazyr (n = 1, SCI), Haegarda (n = 1, SCI), Takhzyro (n = 1, SCI), Beriner (n = 3, IVI), Ruconest (n = 3), and Orladeyo (n = 1, pill, off-label use).

^c Ten adults took Firazyr (n = 8, SCI), Haegarda (n = 1, SCI), and Beriner (n = 1, IVI).

^d Multiple response question; percentages sum to greater than 100% per column.

Conclusions

- All participants preferred the hypothetical oral on-demand treatment over hypothetical self-administered injection on-demand treatment when efficacy and tolerability/mild side-effect risk were similar
- Effectiveness was a commonly reported 'like' and injection-site pain/burning was a commonly reported 'dislike' of current on-demand treatment
- In the hypothetical comparison, self-administered injection was only preferred over oral on-demand treatment if it offered substantially better efficacy over oral treatment, and only if the oral treatment had substantively worse tolerability/side-effect risk than observed in available clinical studies
- Quantitative analyses in a larger cohort are warranted to better refine on-demand treatment preferences, for shared decision-making

References

1. Bork K, Anderson JT, Caballero T, Craig T, Johnston DT, Li HH, et al. Assessment and management of disease burden and quality of life in patients with hereditary angioedema: a consensus report. Allergy Asthma Clin Immunol. 2021; 17: 40

Presented: 2023 US HAEA National Summit, July 20 - 23, 2023 in Orlando, Florida.

Please scan this QR code to view the poster after the congress.

